

RESEARCH

Open Access



Free-living gait does not differentiate chronic mTBI patients compared to healthy controls

Dylan Powell¹, Alan Godfrey¹, Lucy Parrington^{2,3}, Kody R. Campbell², Laurie A. King² and Sam Stuart^{2,4,5*} 

Abstract

Background: Physical function remains a crucial component of mild traumatic brain injury (mTBI) assessment and recovery. Traditional approaches to assess mTBI lack sensitivity to detect subtle deficits post-injury, which can impact a patient's quality of life, daily function and can lead to chronic issues. Inertial measurement units (IMU) provide an opportunity for objective assessment of physical function and can be used in any environment. A single waist worn IMU has the potential to provide broad/macro *quantity* characteristics to estimate gait mobility, as well as more high-resolution micro spatial or temporal gait characteristics (herein, we refer to these as measures of *quality*). Our recent work showed that quantity measures of mobility were less sensitive than measures of turning quality when comparing the free-living physical function of chronic mTBI patients and healthy controls. However, no studies have examined whether measures of gait quality in free-living conditions can differentiate chronic mTBI patients and healthy controls. This study aimed to determine whether measures of free-living gait quality can differentiate chronic mTBI patients from controls.

Methods: Thirty-two patients with chronic self-reported balance symptoms after mTBI (age: 40.88 ± 11.78 years, median days post-injury: 440.68 days) and 23 healthy controls (age: 48.56 ± 22.56 years) were assessed for ~7 days using a single IMU at the waist on a belt. Free-living gait quality metrics were evaluated for chronic mTBI patients and controls using multi-variate analysis. Receiver operating characteristics (ROC) and Area Under the Curve (AUC) analysis were used to determine outcome sensitivity to chronic mTBI.

Results: Free-living gait quality metrics were not different between chronic mTBI patients and controls (all $p > 0.05$) whilst controlling for age and sex. ROC and AUC analysis showed stride length (0.63) was the most sensitive measure for differentiating chronic mTBI patients from controls.

Conclusions: Our results show that gait quality metrics determined through a free-living assessment were not significantly different between chronic mTBI patients and controls. These results suggest that measures of free-living gait quality were not impaired in our chronic mTBI patients, and/or, that the metrics chosen were not sensitive enough to detect subtle impairments in our sample.

Keywords: mTBI, Concussion, Inertial measurement unit, Gait

Introduction

Traumatic brain injuries (TBI) can be broadly defined as sudden trauma causing damage to the brain, with severity ranging from mild TBI (mTBI; commonly known as concussion) to severe TBI [1]. An array of impairments accompany TBI, such as deficits in physical (balance, gait

*Correspondence: sam.stuart@northumbria.ac.uk

² Department of Neurology, Oregon Health and Science University, Portland, OR, USA

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

and turning) [2, 3], psychological (cognitive impairments and symptoms) [4], and sensory function (visual or vestibular deficits) [5]. Such deficits can be subtle and difficult to detect in mTBI and may persist for long periods after the initial injury (e.g., >3 months). Chronic symptoms post-mTBI can significantly impact quality of life and daily function, which can lead to prolonged issues/symptoms [6]. Physical impairments are especially prevalent in mTBI, with eight out of ten people with acute mTBI reporting balance impairments within a few days of the injury and three out of ten reporting longer-term (chronic) balance or gait impairments [5, 7, 8]. Therefore, physical testing (balance and gait) remains a crucial component of clinical assessment to quantify impairment across various mTBI timelines [9–12]. Understanding gait and balance deficits may provide targets for rehabilitation.

Balance impairment is commonly assessed in the acute stage following mTBI [13, 14], primarily using the Balance Error Scoring System (BESS). The BESS requires a clinician to manually record errors each time the patient fails to maintain a balance stance position. However, the sensitivity of the BESS is highly variable due to considerable subjectivity in error counting, which impacts the replicability and validity of results [15–18]. Additionally, subtle balance deficits may be visually undetectable by a clinician's subjective assessment and therefore unmeasurable. Other physical impairments, such as gait deficits, are often not examined by clinicians following acute mTBI. Tandem gait/walking may be done as part of the Sports Concussion Assessment Tool (SCAT), however clinician observation has been found to miss subtle gait deficits that persist in chronic mTBI patients (i.e. due to low ceiling effect of the test) [19]. To detect subtle gait deficits following mTBI, assessment is typically conducted in research settings with objective laboratory equipment, such as force plates and 3D motion capture [7, 20–23]. As such, there have been improvements in objective and instrumented assessment which can yield greater sensitivity than traditional qualitative methods of assessment [14].

Results from laboratory-based objective gait assessment have found pace-related deficits (stride length and gait speed) in chronic mTBI patients compared with healthy controls [24], suggesting gait may be a useful diagnostic marker of mTBI. While laboratory studies provide a foundation for evaluating the differences between healthy and impaired gait, laboratory-centric assessment methods are prescriptive in nature, and may mask subtle mTBI-related deficits that may otherwise occur within habitual (free-living) environments. Accordingly, monitoring gait beyond the laboratory may provide an

opportunity to detect subtle and meaningful deficits following mTBI.

Continuous gait monitoring in free-living environments is becoming more common, due to the widespread use of discrete inertial-based measurement units (IMU), which are the accepted standard for gathering continuous, high-resolution data [25, 26]. IMUs can estimate general mobility outcomes (e.g. measures of quantity such as steps per day) or more refined balance, gait and turning outcomes characterising quality of movement within any environment (e.g. stride length or turning speeds) [2, 14, 27–30]. Our recent work examined free-living mobility quantity and turning quality measures in chronic mTBI patients and controls. We found turning quality metrics to be more sensitive than mobility quantity metrics to differentiate groups [3]. Specifically, those with chronic mTBI had larger, slower and more variable turns during daily life, but had a similar number of steps per day compared with controls [3]. While that study evaluated turning quality, it did not measure other gait quality metrics such as stride velocity, step length, or swing time. Additionally, while previous studies have examined mTBI gait in research settings, no study to date has comprehensively quantified free-living gait quality in chronic mTBI patients and healthy controls. Therefore, a gap remains as to whether measures of free-living gait quality are impaired in chronic mTBI patients. Greater understanding of how mobility is affected in free-living environments may uncover useful markers for subtle deficits in chronic mTBI patients.

The aims of this study were therefore to; (1) explore if free-living gait is impaired in people with chronic mTBI compared with healthy controls, and (2) determine the most sensitive free-living gait quality metrics that differentiate chronic mTBI patients from controls. We hypothesise that free-living mobility would be impaired in chronic mTBI patients compared to controls, with selective gait quality characteristics sensitive to differentiate chronic mTBI.

Methods

Participants

Thirty-two symptomatic chronic mTBI patients and 23 healthy controls participated. Participants were recruited as part of a larger study [31], through posters in athletic facilities, physical therapy clinics, hospitals, concussion clinics, community notice boards, and cafes in and around the Portland, OR metropolitan area. Patient demographics are shown in Table 1. Ethical approval was granted by the Oregon Health and Science University (OHSU) and Veterans Affairs Portland Health Care System (VAPORHCS) joint institutional review board with

Table 1 Participant demographics

	Controls (n = 23)	mTBI (n = 32)	p
Age (years)	48.56 (22.56)	40.88 (11.78)	0.11
Sex (male or female) ^b	M(6) F(17)	M(6) F(26)	0.52
Height (cm)	165.46 (8.03)	168.51 (9.19)	0.22
Mass (kg)	68.03 (15.32)	76.17 (18.80)	0.25
NSI total score	-	35.88 (13.9)	-
NSI vestibular	-	5.44 (2.22)	-
NSI somatosensory	-	10 (4.92)	-
NSI cognitive score	-	8.34 (3.89)	-
NSI affective score	-	10.34 (5.64)	-
Days since injury ^a	-	440.68 (700.63)	-

^a Median and interquartile range

^b chi-squared, Mean and standard deviation reported unless otherwise stated. mTBI, mild traumatic brain injury; NSI—neurobehavioral symptom inventory

participants providing written informed consent before commencing the study.

Inclusion and exclusion criteria

Participants were included in the chronic mTBI group if they had had a diagnosis of mTBI based upon Veteran Health Administration (VHA)/Department of Defense (DoD) [32] criteria and who were greater than three months post mTBI with self-reported balance impairments. The control group consisted of those who had no history of brain injury in the last year. Additionally, mTBI patients were required to have minimal to no cognitive deficits as determined by the Short-Blessed Test (score ≤ 8) [33] and no peripheral vestibular or oculomotor pathology preceding their mTBI. Participants were excluded if they had any musculoskeletal injury which could impair their gait or balance or a recent history of moderate or severe substance abuse.

Gait analysis

Participants were asked to wear an IMU for 7 days, and participants with less than 3 days were excluded from analysis, in line with previous studies [3, 34, 35]. Participants wore a compact (L \times W \times H: 43.7 \times 39.7 \times 13.7 mm, 128 Hz) and lightweight (<25 g) IMU (previously validated [36–38]) attached to a belt (128 Hz, Opal V1, APDM Inc., Portland, OR) that contained an accelerometer (± 16 g, ± 200 g) and gyroscope (± 2000 deg/s). Participants wore the IMU around their waist for a minimum of 5 h per day for up to 7 days using the protocol described previously by Fino et al. 2017 [31] and Stuart et al. 2020 [3]. Data were stored on the IMU internal storage (8 Gb) and then downloaded via proprietary software (MobilityLab, APDM Inc., Portland, OR) to a laptop.

Free-living data were then processed using custom-made and validated MATLAB[®] (MathWorks Inc, Massachusetts, USA) algorithms to estimate 12 free-living gait quality metrics [34, 35, 39, 40].

Gait

Free-living measures of gait quality were calculated using a bespoke MATLAB[®] algorithm as follows. The waist worn IMU was used to examine orientation and periods of static and dynamic activity [39, 40]. Subsequently, the latter were examined for initial and final foot contact events within the gait cycle via the continuous wavelet transform [41], where a bout/period of walking was predefined by a time period of between 0.25 and 2.25 s and ≥ 3 steps [42]. For the purposes of this study, a movement bout was classified as > 10 s. Gait quality metrics included mean; stance time (seconds, s), step time(s), stride time (s), swing time (s), stride length (centimetres, cm), stride velocity (cm/second, cms^{-1}) and coefficient of variation (CV) of these measures.

Self-reported symptoms

Chronic mTBI patients completed the Neurobehavioral Symptom Inventory (NSI) which is widely used in the assessment of mTBI symptoms [24, 43]. The NSI is composed of 22 items within the questionnaire and recorded on a five-point Likert scale, with higher scores indicating more severe symptoms. The maximum a participant can score is 88. The NSI and subscales [44] have acceptable reliability in characterising presence and tracking severity of symptoms in TBI [44, 45]. The NSI remains the cornerstone of clinical symptom assessment and was determined as the appropriate method to capture self-reported impairments in the chronic mTBI patients.

Statistical analysis

Data were analysed in SPSS (v23, IBM) and R studio (Boston, MA, USA). All data were normally distributed as assessed with Shapiro-Wilks tests and therefore parametric tests were used. Independent t-tests were performed comparing demographic information between mTBI and control groups. To compare free-living gait quality metrics between chronic mTBI patients and controls, we used separate multivariate analysis of covariance (MANCOVA). MANCOVA was used to control for sex and age [4, 46].

To estimate which gait quality metrics differentiated chronic mTBI patients from controls, we used receiver operating characteristic (ROC) and area under the curve (AUC) analysis. ROC analysis provides a trade-off between specificity and sensitivity between the various free-living gait quality metrics and binary classification of either mTBI patients and healthy control. Statistical

significance was determined at $p < 0.05$ (*two-tailed*) unless otherwise stated. Bonferroni corrected significance values were applied for multiple comparisons in free-living gait quality measures ($p < 0.002$). Effect sizes were interpreted as small (0.01), medium (0.06), and large (0.14) as previously described [47].

Results

Demographics and clinical assessments

Demographic characteristics are presented in Table 1 for age (years), height (cm), mass (kg) and the number of days since injury and NSI for the mTBI group only. In our mTBI cohort, NSI total score was moderately high (5th to 9th percentile) compared to previously published normative mTBI scores, demonstrating that our chronic mTBI group was still symptomatic at least more than 3 months after injury [44].

Adherence to IMU device

Participants were asked to wear the IMU-based device for 7 days, but compliance was variable across both groups with several mTBI ($n = 16$) and control ($n = 13$) participants wearing the sensor for less than 7 days. Specifically, the mean number of days that the IMU was worn was 6.8 (± 2.4) days in the mTBI group and 6.04 (± 2.0) days in the control group.

Group differences in free-living gait quality measures

When controlling for age and sex, there were no significant differences in measures of free-living gait quality between chronic mTBI patients ($p > 0.05$) and controls.

Descriptive data for free-living gait quality metrics are provided in Table 2.

Sensitivity and specificity of free-living gait metrics

Figure 1 shows the receiver operating characteristics (ROC) analysis for the top four gait quality metrics ($AUC > 0.51$). Free-living gait quality (mean AUC : 0.51) was considered poor at differentiating chronic mTBI patients from controls ($AUC > 0.50$, Table 2).


Discussion

This study progresses our previous work [3], which examined free-living activity quantity and turning quality measured by a single IMU in those with chronic mTBI compared to healthy controls. Free-living mobility assessment in mTBI is still an emerging research area, but results from other neurological conditions (e.g. Parkinson's disease) suggest that impaired gait occurs in parallel with neurological dysfunction [48]. However, results in this study indicated that free-living gait quality was not significantly different between our samples of chronic mTBI patients and healthy controls (when controlling for age and gender). The absence of significant differences in this study are likely multifactorial and could involve both inherent limitations of self-reporting of balance issues, and the chronicity of this mTBI cohort. However, assessment of free-living mobility in chronic mTBI may still allow for improved diagnostics and monitoring of recovery within real-world environments, which is unachievable using analog (non-digital) approaches or laboratory-based assessments only, but further research

Table 2 Free-living gait quality metrics; group differences whilst controlling for age and sex, Area under the Curve (AUC)

Free-living gait metric	mTBI ($n = 32$) mean (S.D.)	Controls ($n = 23$) mean (S.D.)	F	p	η_p^2	AUC
Mean stance time (seconds, s)	0.83 (0.05)	0.85 (0.09)	0.19	0.66	0.00	0.44
Mean step time (s)	0.70 (0.05)	0.73 (0.09)	0.21	0.65	0.00	0.44
Mean stride time (s)	1.41 (0.10)	1.45 (0.18)	0.21	0.65	0.00	0.44
Mean swing time (s)	0.58 (0.05)	0.60 (0.09)	0.22	0.64	0.00	0.44
Mean stride length (centimetres, cm)	74.01 (4.10)	72.68 (3.60)	2.84	0.10	0.05	0.63
Mean stride velocity (cms^{-1})	105.59 (8.88)	101.34 (11.47)	1.37	0.25	0.03	0.60
Stance time variability CV (s)	0.20 (0.01)	0.21 (0.02)	0.03	0.87	0.00	0.49
Step time variability CV (s)	0.20 (0.01)	0.20 (0.02)	0.10	0.75	0.00	0.48
Stride time variability CV (s)	0.22 (0.01)	0.22 (0.01)	0.35	0.56	0.01	0.51
Swing time variability CV (s)	0.20 (0.01)	0.21 (0.02)	0.13	0.72	0.00	0.47
Step length variability CV (s)	18.62 (1.18)	18.32 (0.96)	2.30	0.14	0.04	0.61
Step velocity variability CV (cms^{-1})	36.90 (3.11)	35.48 (4.08)	1.18	0.28	0.02	0.60

Bolded p values; $p < 0.05$ (Bonferroni corrected p value 0.002). Group analysis of covariance results controlling for age and sex. mTBI, mild traumatic brain injury; S.D., standard deviation; CV, coefficient of variation, η_p^2 partial eta squared of effect size, F Wilks' λ , $AUC > 0.50$ in italics and bold



with longitudinal assessments following the initial injury would be required.

Free-living gait quality measures are not impaired in chronic mTBI patients

Our results show that free-living gait quality metrics were not different between chronic mTBI and control groups, which is surprising given this cohort had self-reported balance deficits. Overall research into chronic mTBI has yet to gain consensus on what specific measures can differentiate healthy people from those with mTBI [24]. Indeed some laboratory-based studies have found pace-related deficits (stride length and gait speed) while other studies have found no differences outside of the acute timeframe (>10 days) [2]. Laboratory gait assessment does allow for more controlled assessment of complex tasks (e.g. dual-task, obstacle avoidance, etc.), which may be required to elicit or provoke gait deficits in chronic mTBI [2, 49]. For example, dual-task laboratory assessment in people with chronic mTBI can reveal gait deficits in rhythm (stride time) [24]. However, complex laboratory tasks fail to fully replicate free-living environments where motor, cognitive and sensory function are continuously challenged [50]. Given these challenges in free-living environments, we were surprised that our

measures of gait quality did not suggest impaired mobility in this chronic mTBI cohort.

The lack of significant differences and low effect sizes in gait quality measures between chronic mTBI patients and healthy controls may be related to the considerable chronicity (median 1.2 years post-injury) of this mTBI cohort. This duration may have resulted in the cohort developing chronic compensatory strategies over time to replicate 'normal' gait patterns during walking in their daily life. To fully understand this, future research should test participants in both the laboratory under complex conditions (e.g., dual-task, obstacle walking, turns course etc.) and in free-living environments longitudinally from the time of initial injury to better understand how gait changes acutely after mTBI and into more chronic stages. Similarly incorporating assessment of turning, which is a more complex task that is difficult to compensate for, may also reveal subtle mobility deficits [24, 28, 51]. Overall, there is no definitive way of objectively understanding the reasons for lack of differences in free-living gait quality between our cohorts of chronic mTBI patients and healthy controls. There are many unknown factors and contexts that affect free-living assessments. For example, here the environments participants were regularly walking in, the surfaces they walked on, or the types of terrain encountered were all unknown and such heterogeneity

could impact results [52]. Equally, it is not possible to quantify the usual free-living mobility habits of the participants or to determine if this chronic mTBI cohort displayed any compensatory behaviour strategies (e.g., refraining from talking or performing other tasks whilst walking) that could further impact results. The introduction of egocentric video recordings of free-living mobility may enable greater insight and a robust reference to better understand the context of environments [53]. If used in conjunction with objective free-living IMU assessment, video data could yield even greater contextual understanding of free-living gait performance and any compensatory behaviour mTBI patients display within an environment.

Strengths and limitations

Digital technologies such as IMU's have many advantages over traditional methods of assessment including objectivity and continuous data collection. The primary strength of this study was the use of a single IMU to objectively measure free-living gait quality in chronic mTBI patients and controls; the use of a single device and assessment within usual daily life means that subjects had low research burden [54]. We also quantified useful gait quality metrics from clinical-based conceptual models from neurological-based research. Although use of a single IMU alone on the lower back facilitated more rapid data collection and reduced burden, it fails to quantify other useful gait characteristics which may provide more insight to dynamic postural control and environmental information i.e., step width and step width variability arising from uneven terrain [55]. Thus, future research should investigate additional gait characteristics (based on conceptual gait models) with e.g., multiple IMU's (on the feet) or a video-based wearable for a more informed free-living assessment. While the authors are not currently aware of any IMU-based technology to quantify step width during free-living, a computer vision approach has been suggested from a wearable camera [53]. Additionally, the outcome measures presented are primarily research-orientated, requiring a great deal of time-consuming post-processing and checking, which is based on prior experience of inertial data [56, 57]. Therefore, there are needs to refine and deploy software that clinicians and patients can easily navigate, which would allow more widespread uptake and use by health professionals [57].

No power calculation was used in this study as it was based as an exploratory study with opportunistic sampling. This may have limited the strength of any conclusions drawn and should be taken with caution. Future research should aim to utilise power calculations to ensure sufficient sample size and ability to detect

small differences in results. Participants were assessed for ~7 days using a single IMU attached to a waist belt. However variation in the exact length of time participants wore wearables (minimum three days) could introduce differences and therefore not reflect true habitual free-living mobility as used in other studies [48, 58]. Using multiple IMUs may provide more detailed spatial and temporal data for turning, balance and gait as used in previous studies [24], but this carries different limitations; such as longer data download, processing complexity and increased wearer burden, limiting the practical or clinical application. This trade-off should be considered in future studies as a potential improvement to the assessment protocol. [59, 60].

There were some additional limitations to this study. First, a more detailed demographic profile could be reported in future studies to derive further inferences about the free-living mobility results or underlying physiological mechanisms for persistent symptom and mobility deficits [24]. For example, the symptom questionnaires were limited to NSI that were only completed by the mTBI cohort, which limited any useful comparisons and inference on the relationship between groups [3]. Second, balance problems in the chronic mTBI group were self-reported with no baseline or robust analysis done to quantify the magnitude of impairment [3], with the many factors such as the previous history of mTBI and evidence of abnormal neuroimaging omitted [4, 61]. Third, the differences in this mTBI cohort's chronicity are likely to limit the direct comparison with other studies. Our study's cohort was chronic with a median post-injury time greater than 1-year, which compared to other studies examining people post-mTBI is a longer time since injury [24, 62].

Conclusions

Our results demonstrate that free-living IMU-based gait quality metrics were not significantly different between patients with chronic mTBI and healthy aged-matched controls. Despite a lack of significant findings herein, we feel that there is value in undertaking free-living mobility assessments. This study has highlighted that a single IMU can obtain a wealth of continuous free-living gait quality measures in people with symptomatic chronic mTBI and healthy controls. While this exploratory study indicated no between group differences, we feel that this work provides a foundation for future work in this area, where a-priori power and sample size are controlled. When considering the results of this study with our previous findings [3], we advocate that assessments of free-living mobility should include both measures of gait and turning quality. Future research should also focus on (i) additional gait characteristics from conceptual gait models

and (ii) longitudinal analysis of chronic mTBI patients during different stages of recovery (acute to chronic) to holistically monitor mobility impairments and recovery. Improving objectivity in mTBI assessment will result in greater understanding of injury progression, recovery, and rehabilitation across a variety of clinical settings.

Acknowledgements

The authors would like to thank all those who volunteered for the study.

Author contributions

DP was responsible for writing the manuscript, editing, statistical and data analysis. AG was responsible for data analysis, editing and reviewing the manuscript. LP, KRC and LAK were responsible for data collection and reviewing the manuscript. SS was responsible for data analysis, writing, statistical analysis and reviewing the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by the Assistant Secretary of Defense for Health Affairs under Award No. W81XWH-15-1-0620. Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the Department of Defense. D Powell is supported by a Northumbria University doctoral research programme and the Private Physiotherapy Education Foundation (PI Powell: PPEF#349, RPJ03732) and (PI Stuart PPEF#368). Dr Stuart is supported by research funding from the Parkinson's Foundation via a post-doctoral fellowship and clinical research award (PF-FBS-1898, PF-CRA-2073).

Availability of data and materials

De-identified data generated from this study will be deposited into the Federal Interagency Traumatic Brain Injury Research (FITBIR) Informatics System.

Declarations

Ethics approval and consent to participate

Ethical approval was granted by the Oregon Health and Science University (OHSU) and Veterans Affairs Portland Health Care System (VAPORHCS) joint institutional review board with participants providing written informed consent before commencing the study.

Consent for publication

Not applicable.

Competing interests

All authors declare that they have no competing interests.

Author details

¹Department of Computer and Information Sciences, Northumbria University, Newcastle-upon-Tyne, UK. ²Department of Neurology, Oregon Health and Science University, Portland, OR, USA. ³Department of Dietetics, Human Nutrition and Sport, La Trobe University, Victoria, Australia. ⁴Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle-upon-Tyne NE1 8ST, UK. ⁵North Tyneside Hospital, Northumbria Healthcare NHS Foundation Trust, North Shields, UK.

Received: 21 July 2021 Accepted: 18 May 2022

Published online: 26 May 2022

References

- National Institute of Neurological Disorders. Traumatic Brain Injury Information Page|National Institute of Neurological Disorders and Stroke. 2019. <https://www.ninds.nih.gov/Disorders/All-Disorders/Traumatic-Brain-Injury-Information-Page>
- Fino PC, Parrington L, Pitt W, Martini DN, Chesnutt JC, Chou LS, et al. Detecting gait abnormalities after concussion or mild traumatic brain injury: a systematic review of single-task, dual-task, and complex gait. *Gait Posture*. 2018;157–66
- Stuart S, Parrington L, Martini DN, Kreter N, Chesnutt JC, Fino PC, et al. Analysis of free-living mobility in people with mild traumatic brain injury and healthy controls: quality over quantity. *J Neurotrauma*. 2020;37:139–45. <https://doi.org/10.1089/neu.2019.6450>
- Kenzie ES, Parks EL, Bigler ED, Wright DW, Lim MM, Chesnutt JC, et al. The dynamics of concussion: Mapping pathophysiology, persistence, and recovery with causal-loop diagramming. *Front Neurol*. 2018;9:1.
- Alsalaheen BA, Mucha A, Morris LO, Whitney SL, Furman JM, Camiollo-Reddy CE, et al. Vestibular rehabilitation for dizziness and balance disorders after concussion. *J Neurol Phys Ther*. 2010;87–93
- Chu SY, Tsai YH, Xiao SH, Huang SJ, Yang CC. Quality of return to work in patients with mild traumatic brain injury: a prospective investigation of associations among post-concussion symptoms, neuropsychological functions, working status and stability. *Brain Inj*. 2017;31:1674–82.
- Basford JR, Chou LS, Kaufman KR, Brey RH, Walker A, Malec JF, et al. An assessment of gait and balance deficits after traumatic brain injury. *Arch Phys Med Rehabil*. 2003;84:343–9.
- Kleffeldgaard I, Roe C, Soberg HL, Bergland A. Associations among self-reported balance problems, post-concussion symptoms and performance-based tests: a longitudinal follow-up study. *Disabil Rehabil*. 2012;34:788–94.
- Godfrey A, Lara J, Del Din S, Hickey A, Munro CA, Wuiff C, et al. iCap: instrumented assessment of physical capability. *Maturitas*. 2015;82:116–22.
- Alberts JL, Hirsch JR, Koop MM, Schindler DD, Kana DE, Linder SM, et al. Using accelerometer and gyroscopic measures to quantify postural stability. *J Athl Train*. 2015;50:578–88.
- McCrorry P, Meeuwisse W, Dvorak J, Aubry M, Bailes J, Broglio S, et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. *Br J Sport Med*. 2017;12:6–11. <https://doi.org/10.1136/bjsports-2017-097699>
- Stuart S, Hickey A, Morris R, O'Donovan K, Godfrey A. Concussion in contact sport: a challenging area to tackle. *J Sport Heal Sci*. 2017;6:299–301.
- Bell DR, Guskiewicz KM, Clark MA, Padua DA. Systematic review of the balance error scoring system. *Sports Health*. 2011;287–95.
- King LA, Mancini M, Fino PC, Chesnutt J, Swanson CW, Markwardt S, et al. Sensor-based balance measures outperform modified balance error scoring system in identifying acute concussion. *Ann Biomed Eng*. 2017;45:2135–45.
- Cooper R, Kuh D, Hardy R. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *BMJ*. 2010;341:639.
- Cooper R, Kuh D, Cooper C, Gale CR, Lawlor DA, Matthews F, et al. Objective measures of physical capability and subsequent health: a systematic review. *Age Ageing* 2011;14–23
- Finnoff JT, Peterson VJ, Hollman JH, Smith J. Intrarater and interrater reliability of the balance error scoring system (BESS). *PM R*. 2009;1:50–4.
- Powell D, Stuart S, Godfrey A. Sports related concussion: an emerging era in digital sports technology. *NPJ Digit Med*. 2021;4:1–8.
- Howell DR, Osternig LR, Chou LS. Single-task and dual-task tandem gait test performance after concussion. *J Sci Med Sport*. 2017;20:622–6.
- Chesnutt J, King L, Parrington L, Fino P, Stuart S. Turning the tide: real-world turns are more sensitive to mTBI deficits than daily activity measures. *Arch Phys Med Rehabil*. 2018;99:e181.
- King LA, Horak FB, Mancini M, Pierce D, Priest KC, Chesnutt J, et al. Instrumenting the balance error scoring system for use with patients reporting persistent balance problems after mild traumatic brain injury. *Arch Phys Med Rehabil*. 2014;95:353–9. <https://doi.org/10.1016/j.apmr.2013.10.015>
- Lee H, Sullivan SJ, Schneiders AG. The use of the dual-task paradigm in detecting gait performance deficits following a sports-related concussion: a systematic review and meta-analysis. *J Sci Med Sport* 2013;2–7
- Schneiders AG, Sullivan SJ, Handcock P, Gray A, McCrorry PR. Sports concussion assessment: the effect of exercise on dynamic and static balance. *Scand J Med Sci Sport*. 2012;22:85–90.
- Martini DN, Parrington L, Stuart S, Fino PC, King LA. Gait performance in people with symptomatic, chronic mild traumatic brain injury. *J Neurotrauma*. 2021;38:218–24. <https://doi.org/10.1089/neu.2020.6986>

25. Seshadri DR, Li RT, Voos JE, Rowbottom JR, Alfes CM, Zorman CA, et al. Wearable sensors for monitoring the internal and external workload of the athlete. *NPJ Digit Med*. 2019;2.
26. Powell D, Stuart S, Godfrey A. Wearables in rugby union: a protocol for multimodal digital sports-related concussion assessment. *PLoS ONE*. 2021;16:e0261616. <https://doi.org/10.1371/journal.pone.0261616>.
27. Pitt W, Chen SH, Chou LS. Using IMU-based kinematic markers to monitor dual-task gait balance control recovery in acutely concussed individuals. *Clin Biomech*. 2020;80:105145.
28. Stuart S, Parrington L, Morris R, Martini DN, Fino PC, King LA. Gait measurement in chronic mild traumatic brain injury: a model approach. *Hum Mov Sci* 2020;69
29. Howell D, Osternig L, Chou L-SS. Monitoring recovery of gait balance control following concussion using an accelerometer. *J Biomech*. 2015;48:3364–8.
30. Del Din S, Godfrey A, Galna B, Lord S, Rochester L. Free-living gait characteristics in ageing and Parkinson's disease: impact of environment and ambulatory bout length. *J Neuroeng Rehabil*. 2016;13:1–12. <https://doi.org/10.1186/s12984-016-0154-5>.
31. Fino PC, Peterka RJ, Hullar TE, Murchison C, Horak FB, Chesnutt JC, et al. Assessment and rehabilitation of central sensory impairments for balance in mTBI using auditory biofeedback: a randomized clinical trial. *BMC Neurol*. 2017;17:41. <https://doi.org/10.1186/s12883-017-0812-7>.
32. Veterans Affairs/Department of Defense. Management of concussion-mild traumatic brain injury (mTBI) clinical practice guidelines. 2016. <https://www.healthquality.va.gov/guidelines/rehab/mtbi/>
33. Katzman R, Brown T, Fuld P, Peck A, Schechter R, Schimmel H. Validation of a short orientation-memory-concentration test of congestive impairment. *Am J Psychiatry*. 1983;140:734–9.
34. Mancini M, El-Gohary M, Pearson S, McNames J, Schlueter H, Nutt JG, et al. Continuous monitoring of turning in Parkinson's disease: rehabilitation potential. *NeuroRehabilitation*. 2015;37:3–10.
35. El-Gohary M, Pearson S, McNames J, Mancini M, Horak F, Mellone S, et al. Continuous monitoring of turning in patients with movement disability. *Sensors (Switzerland)*. 2014;14:356–69.
36. Washabaugh EP, Kalyanaraman T, Adamczyk PG, Claflin ES, Krishnan C. Validity and repeatability of inertial measurement units for measuring gait parameters. *Gait Posture*. 2017;55:87–93.
37. Taylor L, Miller E, Kaufman KR. Static and dynamic validation of inertial measurement units. *Gait Posture*. 2017;57:80–4.
38. Schmitz-Hübisch T, Brandt AU, Pfueller C, Zange L, Seidel A, Kühn AA, et al. Accuracy and repeatability of two methods of gait analysis—GaitRite™ und mobility Lab™—in subjects with cerebellar ataxia. *Gait Posture*. 2016;48:194–201.
39. Hickey A, Del Din S, Rochester L, Godfrey A. Detecting free-living steps and walking bouts: validating an algorithm for macro gait analysis. *Physiol Meas*. 2017;38:N1–15. <https://doi.org/10.1088/1361-6579/38/1/N1>.
40. Godfrey A, Del Din S, Barry G, Mathers JC, Rochester L. Instrumenting gait with an accelerometer: a system and algorithm examination. *Med Eng Phys*. 2015;37:400–7.
41. McCamley J, Donati M, Grimpampi E, Mazzà C. An enhanced estimate of initial contact and final contact instants of time using lower trunk inertial sensor data. *Gait Posture*. 2012;36:316–8.
42. Najafi B, Aminian K, Paraschiv-Ionescu A, Loew F, Büla CJ, Robert P. Ambulatory system for human motion analysis using a kinematic sensor: Monitoring of daily physical activity in the elderly. *IEEE Trans Biomed Eng*. 2003;50:711–23.
43. Belanger HG, Silva MA, Donnell AJ, McKenzie-Hartman T, Lamberty GJ, Vanderploeg RD. Utility of the neurobehavioral symptom inventory as an outcome measure: a VA TBI model systems study. *J Head Trauma Rehabil*. 2017;32:46–54.
44. Vanderploeg RD, Silva MA, Soble JR, Curtiss G, Belanger HG, Donnell AJ, et al. The structure of postconcussion symptoms on the neurobehavioral symptom inventory: a comparison of alternative models. *J Head Trauma Rehabil*. 2015;30:1–11.
45. Black AM, Miutz LN, Warriyar VK, Schneider KJ, Owen Yeates K, Emery CA. Baseline performance of high school rugby players on the sport concussion assessment tool 5. *J Athl Train*. 2020;55:116–23.
46. Anto-Ocrah M, Tiffany K, Hasman L, Van Wijngaarden E. Mild traumatic brain injury/concussion and female sexuality, a scoping review of the literature. *Inj Epidemiol*. 2020;7:1. <https://doi.org/10.1186/s40621-020-0232-9>.
47. Kontos AP, Eagle SR, Mucha A, Kochick V, Reichard J, Moldovan C, et al. A randomized controlled trial of precision vestibular rehabilitation in adolescents following concussion: preliminary findings. *J Pediatr*. 2021;239:193–9.
48. Del Din S, Galna B, Godfrey A, Bekkers EMJ, Pelosin E, Nieuwhof F, et al. Analysis of free-living gait in older adults with and without Parkinson's disease and with and without a history of falls: identifying generic and disease-specific characteristics. *Journals Gerontol - Ser A Biol Sci Med Sci*. 2019;74:500–6.
49. Parrington L, Fino PC, Swanson CW, Murchison CF, Chesnutt J, King LA. Longitudinal assessment of balance and gait after concussion and return to play in collegiate athletes. *J Athl Train*. 2019;54:429–38.
50. Takakusaki K. Functional neuroanatomy for posture and gait control. *J Mov Disord*. 2017;10:1–17.
51. Martini DN, Broglio SP. Long-term effects of sport concussion on cognitive and motor performance: a review. *Int J Psychophysiol*. 2018;132:25–30.
52. Celik Y, Stuart S, Woo WL, Godfrey A. Gait analysis in neurological populations: progression in the use of wearables. *Med. Eng. Phys*. 2021;9–29.
53. Nouredanesh M, Li AW, Godfrey A, Hoey J, Tung J. Chasing feet in the wild: a proposed egocentric motion-aware gait assessment tool. *Lect Notes Comput Sci (including Subser Lect Notes Artif Intell Lect Notes Bioinformatics)*. 2019. https://doi.org/10.1007/978-3-030-11024-6_12.
54. Lord S, Galna B, Rochester L. Moving forward on gait measurement: toward a more refined approach. *Mov Disord*. 2013;28:1534–43. <https://doi.org/10.1002/mds.25545>.
55. Morris R, Hickey A, Del Din S, Godfrey A, Lord S, Rochester L. A model of free-living gait: a factor analysis in Parkinson's disease. *Gait Posture*. 2017;52:68–71.
56. Godfrey A, Goldsack JC, Tenaerts P, Coravos A, Aranda C, Hussain A, et al. BioMeT and algorithm challenges: a proposed digital standardized evaluation framework. *IEEE J Transl Eng Heal Med*. 2020;8.
57. Godfrey A, Vandendriessche B, Bakker JP, Fitzer-Attas C, Gujar N, Hobbs M, et al. Fit-for-purpose biometric monitoring technologies: leveraging the laboratory biomarker experience. *Clin Transl Sci*. 2021. <https://doi.org/10.1111/cts.12865>.
58. Del Din S, Godfrey A, Mazzà C, Lord S, Rochester L. Free-living monitoring of Parkinson's disease: lessons from the field. *Mov Disord*. 2016;31:1293–313. <https://doi.org/10.1002/mds.26718>.
59. Trojaniello D, Cereatti A, Della Croce U. Accuracy, sensitivity and robustness of five different methods for the estimation of gait temporal parameters using a single inertial sensor mounted on the lower trunk. *Gait Posture*. 2014;40:487–92.
60. Goldsack JC, Coravos A, Bakker JP, Bent B, Dowling AV, Fitzer-Attas C, et al. Verification, analytical validation, and clinical validation (V3): the foundation of determining fit-for-purpose for Biometric Monitoring Technologies (BioMeTs). *NPJ Digit Med Nat Res*. 2020;3:1–15. <https://doi.org/10.1038/s41746-020-0260-4>.
61. MacDonald SWS, Nyberg L, Bäckman L. Intra-individual variability in behavior: links to brain structure, neurotransmission and neuronal activity. *Trends Neurosci*. 2006;29:474–80.
62. Fino PC, Parrington L, Walls M, Sippel E, Hullar TE, Chesnutt JC, et al. Abnormal turning and its association with self-reported symptoms in chronic mild traumatic brain injury. *J Neurotrauma*. 2018;35:1167–77.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.